

## **Clinical Policy: Penicillamine (Cuprimine)**

Reference Number: CP.PCH.09

Effective Date: 12.01.18

Last Review Date: 11.24

Line of Business: Commercial, HIM

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Penicillamine (Cuprimine<sup>®</sup>) is a chelating agent.

### **FDA Approved Indication(s)**

Cuprimine is indicated for the treatment of:

- Wilson's disease
- Cystinuria
- Severe, active rheumatoid arthritis (RA) in patients who have failed to respond to an adequate trial of conventional therapy

Limitation(s) of use: Available evidence suggests that Cuprimine is not of value in ankylosing spondylitis.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Cuprimine is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Wilson's Disease** (must meet all):

1. Diagnosis of Wilson's disease;
2. Member must use generic Depen<sup>®</sup>, unless contraindicated or clinically significant adverse effects are experienced;
3. Dose does not exceed both of the following (a and b):
  - a. 2 g per day;
  - b. 8 capsules per day.

##### **Approval duration:**

**HIM** – 6 months

**Commercial** – 12 months or duration of request, whichever is less

##### **B. Cystinuria** (must meet all):

1. Diagnosis of cystinuria;
2. Failure of a urinary alkalinizing agent (e.g., potassium citrate) unless contraindicated or clinically significant adverse effects are experienced;

**CLINICAL POLICY****Penicillamine**

3. Member must use generic Depen, unless contraindicated or clinically significant adverse effects are experienced;
4. Dose does not exceed both of the following (a and b):
  - a. 4 g per day;
  - b. 16 capsules per day.

**Approval duration:****HIM** – 6 months**Commercial** – 12 months or duration of request, whichever is less**C. Rheumatoid Arthritis (must meet all):**

1. Diagnosis of RA;
2. Member meets one of the following (a or b):
  - a. Failure of  $\geq 3$  consecutive months of methotrexate;
  - b. If intolerance or contraindication to methotrexate, failure of  $\geq 3$  consecutive months of sulfasalazine, leflunomide, or hydroxychloroquine unless contraindicated or clinically significant adverse effects are experienced;
3. Member must use generic Depen, unless contraindicated or clinically significant adverse effects are experienced;
4. Dose does not exceed both of the following (a and b):
  - a. Initial therapy: 250 mg (1 capsule) per day for at least the first month;
  - b. Maintenance therapy: 1.5 g (6 capsules) per day.

**Approval duration:****HIM** – 6 months**Commercial** – 12 months or duration of request, whichever is less**D. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial and HIM.PA.33 for health insurance marketplace; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial and HIM.PA.103 for health insurance marketplace; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial and HIM.PA.154 for health insurance marketplace.

**II. Continued Therapy****A. All Indications in Section 1 (must meet all):**

1. Member meets one of the following (a or b):

**CLINICAL POLICY****Penicillamine**

- a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy;
  3. If request is for a dose increase, new dose does not exceed (a, b, or c):
    - a. Wilson's disease (both i and ii):
      - i. 2 g per day;
      - ii. 8 capsules per day;
    - b. RA (both i and ii):
      - i. 1.5 g per day;
      - ii. 6 capsules per day;
    - c. Cystinuria (both i and ii):
      - i. 4 g per day;
      - ii. 16 capsules per day.

**Approval duration:****HIM** – 12 months**Commercial** – 12 months or duration of request, whichever is less**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial and HIM.PA.33 for health insurance marketplace; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial and HIM.PA.103 for health insurance marketplace; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial and HIM.PA.154 for health insurance marketplace.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial and HIM.PA.154 for health insurance marketplace, or evidence of coverage documents;
- B. Ankylosing spondylitis.

**CLINICAL POLICY**  
**Penicillamine**

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

RA: rheumatoid arthritis

*Appendix B: Therapeutic Alternatives*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
penicillamine (Depen)	<p><b>Wilson’s disease</b> 250 mg PO QID; adjust to achieve urinary copper excretion 0.5-1 mg/day</p> <p><b>Cystinuria</b> 250 mg PO QD; increase gradually to 1-2 g/day in 4 divided doses and adjust to achieve target urinary cysteine excretion</p> <p><b>RA</b> 125-250 mg PO QD; increase at 1-3 month intervals by 125-250 mg/day according to response and tolerance (typical maintenance range: 500-750 mg/day) – if no improvement at 1-1.5 g/day after 3-4 months, therapy should be discontinued as a response is unlikely to occur</p>	<p>Wilson’s disease: 2 g/day (750 mg/day if pregnant)</p> <p>Cystinuria: 4 g/day</p> <p>RA: 1.5 g/day</p>
potassium citrate	<p><b>Cystinuria*</b> 60 mEq/day divided into 3 doses; titrate to achieve a urine pH within target range 6-7</p>	See regimen
methotrexate (Rheumatrex <sup>®</sup> )	<p><b>RA</b> 7.5 mg/week PO, SC, or IM or 2.5 mg PO Q12 hr for 3 doses/week</p>	30 mg/week
sulfasalazine (Azulfidine <sup>®</sup> )	<p><b>RA</b> <u>Initial dose:</u> 500 mg to 1,000 mg PO QD for the first week. Increase the daily dose by 500 mg each week up to a maintenance dose of 2 g/day. <u>Maintenance dose:</u> 2 g/day PO in divided doses</p>	3 g/day
leflunomide (Arava <sup>®</sup> )	<p><b>RA</b> <u>Initial dose (for low risk hepatotoxicity or myelosuppression):</u> 100 mg PO QD for 3 days <u>Maintenance dose:</u> 20 mg PO QD</p>	20 mg/day

**CLINICAL POLICY**  
**Penicillamine**

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
hydroxychloroquine (Plaquenil®)	<b>RA*</b> <u>Initial dose:</u> 400 – 600 mg/day PO QD <u>Maintenance dose:</u> 200 – 400 mg/day PO QD	600 mg/day

*Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.*

*\*Off-label*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): history of penicillamine-related aplastic anemia or agranulocytosis, nursing, patients with RA and cystinuria who are pregnant (exceptions can be made for certain patients with cystinuria), patients with RA and history or other evidence of renal insufficiency
- Boxed warning(s): none reported

*Appendix D: General Information*

- Although the prescribing information for Cuprimine does not include an absolute maximum dose for Wilson’s disease, it notes it is seldom necessary to exceed a dose of 2 g/day. In addition, both the American Association for the Study of Liver Diseases and the European Association for the Study of the Liver do not recommend doses higher than 1.5 g/day due to potential for rapid and often irreversible neurological deterioration.
- In cystinuria, initial therapy includes high fluid intake, sodium and protein restriction, and urinary alkalinization. The preferred agent for urinary alkalinization is potassium citrate. Other agents that can be used include potassium bicarbonate, acetazolamide, and sodium bicarbonate or citrate.
- In RA, failure of methotrexate or disease-modifying antirheumatic drugs is defined as  $\leq 50\%$  decrease in swollen joint count,  $\leq 50\%$  decrease in tender joint count, and  $\leq 50\%$  decrease in erythrocyte sedimentation rate (ESR), or  $\leq 50\%$  decrease in C-reactive protein (CRP).
- Examples of positive response include: Wilson’s disease: reduction in 24-hour urinary copper excretion; cystinuria: reduction in urinary cysteine level; RA: improvement in symptoms.

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Cystinuria	1-4 g/day PO in 4 divided doses	4 g/day
Wilson’s disease	750-1,500 mg/day PO in divided doses	2 g/day
RA	125-250 mg PO QD	1.5 g/day

**VI. Product Availability**

Capsule: 250 mg

## CLINICAL POLICY

### Penicillamine

#### VII. References

1. Cuprimine Prescribing Information. Bridgewater, NJ: Bausch Health US, Inc.; October 2020. Available at: <https://pi.bauschhealth.com/globalassets/BHC/PI/Cuprimine-PI.pdf>. Accessed July 19, 2024.
2. Roberts EA and Schilsky ML. AASLD practice guidelines: Diagnosis and treatment of Wilson disease: an update. *Hepatology*. 2008; 47(6): 2089-2111.
3. European Association for the Study of the Liver. EASL clinical practice guidelines: Wilson’s disease. *J Hepatology*. 2012; 56(3): 671-685.
4. Pearle MS, Goldfarb DS, Assimos DG, et al. Medical management of kidney stones: AUA guideline. Published 2014. Available at: [http://www.auanet.org/guidelines/medical-management-of-kidney-stones-\(2014\)](http://www.auanet.org/guidelines/medical-management-of-kidney-stones-(2014)).
5. Goldstein R, Goldfarb DS. Early recognition and management of rare kidney stone disorders. *Urol Nurs*. 2017 Mar-Apr; 37(2): 81–102.
6. Biyani CS and Cartledge JJ. Cystinuria–diagnosis and management. *EAU-EEU Update Series* 4. 2006: 175-183.
7. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2024. Available at: <https://www.clinicalkey.com/pharmacology/>. Accessed July 25, 2024.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
4Q 2020 annual review: no significant changes; references reviewed and updated.	08.11.20	11.20
Clarify redirection is to generic Depen, modify redirect language to state “Member must use generic Depen”.	06.02.21	
4Q 2021 annual review: no significant changes; updated reference for HIM off-label use to HIM.PA.154 (replaces HIM.PHAR.21); references reviewed and updated.	08.27.21	11.21
Revised approval duration for Commercial line of business from length of benefit to 12 months or duration of request, whichever is less	09.27.21	02.22
4Q 2022 annual review: no significant changes; references reviewed and updated. Template changes applied to other diagnoses/indications and continued therapy section.	06.23.22	11.22
4Q 2023 annual review: no significant changes; references reviewed and updated.	06.28.23	11.23
4Q 2024 annual review: no significant changes; updated Appendix B dosing for RA; references reviewed and updated.	07.19.24	11.24

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical

## CLINICAL POLICY

### Penicillamine

policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise

**CLINICAL POLICY****Penicillamine**

published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.